
The Role of Neuroinflammation in Chronic Illness


Keith W. Kelley, Professor Integrative Immunology and Behavior Program



"...international, interdisciplinary journal devoted to investigation of the physiological systems that integrate behavioral and immunological responses."

Why Am I Here at the Research Advisory Committee on Gulf War Veterans' Illness?

Question	<u>Answer</u>
- I enjoy flying through Chicago O'Hare?	No
- I like to ride the Blue Line of the Boston Subway?	No
- I like to eat shell fish?	No
- I have nothing better to do?	Maybe
- I like to have dinner with a former post-doc at Harvard?	Yes
- I have published some stuff on neuroinflammation?	Yes





Summary of Research Advisory Committee on GWI, November 3, 2008

"Since 1994, ..expenditures of \$440 million as GWI research."

"The biological basis for GWI is unknown, and there is no efficacious treatment."

"GWI research effort has yet to provide tangible results... Few treatments have been studied and none have been shown to provide significant benefit for...ill veterans."

"The 200,000 veterans with GWI complain of <u>diffuse pain</u>, headaches, <u>difficulties in learning and memory</u>, <u>mood changes</u> and <u>unrelenting fatigue</u>."

What do I Want you to Remember?

Hypothesis from RAC Report:

"...neurotoxic Gulf War exposures may activate inflammatory processes in the brain and that increased brain levels of proinflammatory cytokines can produce a complex of multiple symptoms similar to GWI."

Take Home Message:

Given the lack of treatments for GWI, the <u>neuroinflammation hypothesis is worth testing</u> because FDA-approved drugs to reduce neuroinflammation and pre-clinical animal research models are available.

Minimal Requirements for Neuroinflammation

- Microglial activation (ramified to ameboid, MHC-II, NFκβ)
- Pro- and anti-inflammatory cytokine synthesis
- Cox-1 & -2, iNOS
- Activation by peripheral immune signals
- No neutrophil or monocyte recruitment
- Neuronal dysfunction that does not require neuronal death
- Behavioral changes

The Experience of Just Being Sick:


Why do we feel sick and behave in a sick way when we are ill?





Sickness behavior as a new target for drug development


Stephen Kent, Rose-Marie Bluthé, Keith W. Kelley and Robert Dantzer


Sickness behavior refers to the nonspecific symptoms (anorexia, depressed activity, loss of interest in usual activities, disappearance of body-care activities) that accompany the response to infection. Increasing evidence suggests that these symptoms are part of an organized defense response to antigenic challenge and that they are mediated by the neural effects of cytokines such as interleukin 1. An understanding of the mechanisms involved in these effects should permit development of new drugs aimed at decreasing sickness or promoting recovery processes. (Kent et al, 1992, TIPS, 13:24)












They are not blind, just blinded from view due to HIPAA!

Fatigue and Deficits in Motor Coordination Induced by LPS are Exacerbated in IL-10 Knockout Mice **Motor Coordination** Exhaustive Fatigue IL-1β mRNA in Cerebellum (Rotarod) (Treadmill) 60% 200 100 250 Derigo de la constance (Seconds) 150 100 50 0 80 Fold change 60 40 20 0 IL-10 WT IL-10 WT IL-10 WT KO KO KO ☐ Saline \square LPS





Major Depressive Episode (DSM-IV)


- · Depressed mood
- Loss of interest / anhedonia
- Decreased concentration
- Worthlessness or guilt
- Suicidal thoughts
- Insomnia / hypersomnia
- Changes in appetite / weight
- Agitation / slowing
- Fatigue / anergia


Sickness Behavior


- ?
- Loss of interest / anhedonia
- Cognitive disturbances
- ?
- ?
- Altered sleep pattern
- Anorexia
- Reduced locomotor activity
- Lethargy














Potential Existing FDA-Approved Approaches for Neuroinflammation in Gulf War Illness

- -Antibiotics (Minocycline)
- -New Drugs for Fibromyalgia (e.g., SNRI milnacipran)
- -Botanicals (turmeric, resveratrol, luteolin)
- HMG-CoA reductase inhibitors (statins)
- Diet Soluble Fiber like Pectin
- IDO Inhibitors now in Phase I clinical trials
- -Exercise

Summary Neuroinflammation in Pre-Clinical Animal Models

- -Reduces Appetite and Motivation
- -Increases Exhaustive Fatigue (Mental Fatigue?)
- -Increases Sensitivity to Pain
- -Causes Deficits in Learning and Memory
- -Induces Depressive-Like Behaviors

What do I Want you to Remember?

Hypothesis from RAC Report:

"...neurotoxic Gulf War exposures may activate inflammatory processes in the brain and that increased brain levels of proinflammatory cytokines can produce a complex of multiple symptoms similar to GWI."

Take Home Message:

Given the lack of treatments for GWI, the <u>neuroinflammation hypothesis is worth testing</u> because FDA-approved drugs to reduce neuroinflammation and pre-clinical animal research models are available.



<u>Urbana team:</u>

Caroline Andre
Robert Dantzer
Jonathan Godbout
Keith W. Kelley
Marc Lawson
Jason O'Connor
Xun Fu
Yun Xia



Bordeaux team:

Nathalie Castanon Lucile Capuron Jacques Lestage Maite Moreau

