# MINDFULNESS MEDITATION FOR CHRONIC LOW BACK PAIN

#### INTRODUCTION

Globally, low back pain is the leading cause of disability[1]. Chronic low back pain (CLBP)—the second leading cause of lost work time—is common, costly, disabling, and often refractory to existing treatments.[2-5] Worldwide, prevalence rates have neared 20% in those age 20 to 59.[6] Approximately 80% of U.S. adults experience low back pain during their lifetime, with 2%-8% developing chronic back pain.[4] Americans spend at least \$50 billion per year on low back pain, with chronic back pain making up at least 90% of the costs.[4]

Individuals with CLBP often need to resort to opioid pain medications to reduce pain and improve function, and CLBP has been the leading noncancer chronic pain condition for which long-term opioids have been prescribed.[7] Despite using potent opioid analgesics, many patients continue to suffer from CLBP, co-occurring mental health problems, and adverse effects of opioids, including overdose and death.[7-12] There is a critical need for safe and effective treatments for CLBP.[7,9,10,13,14]

### MINDFULNESS MEDITATION FOR CHRONIC LOW BACK PAIN: RESEARCH

Patients with CLBP frequently turn to complementary and integrative health (CIH) approaches. Mindfulness-based interventions, which are both popular and safe, are important examples of popular CIH approaches.[5]

Mindfulness-based interventions, including mindfulness meditation (referred to just as "meditation"), help train the mind in nonjudgmental attention to the present moment experiences.[15] They have great potential as therapies for CLBP.[16-20] These approaches are already widely used to improve health[21-23], well-being, and pain[24,25], and they show sustained effects over time.[18,26] Mindfulness-based interventions can positively impact many chronic health problems, including depression and anxiety[27], which are common in patients with CLBP.[3,9,10,16,18,28]

Mindfulness-based interventions can exert biological effects and have been associated with beneficial changes in brain areas involved in adaptive pain, stress, cognitive, and affect regulation.[29,30] Meditation-learned skills can complement those acquired through "traditional" behavioral therapies, such as Cognitive Behavioral Therapy (CBT), which is a part of evidence-based, standard-of-care treatments for CLBP.[31-33] Refer to "Mindful Awareness" and "Power of the Mind" Whole Health modules for additional information.

Although meditation has been used clinically in CLBP for several decades, research evidence on its efficacy in this specific patient population is limited, albeit promising. Initial uncontrolled studies in the 1980s have shown promise of Mindfulness-Based Stress Reduction (MBSR) for refractory chronic pain and reduction in pain medication use.[34,35]

Since then, several small randomized controlled trials have examined mindfulness-based therapy's efficacy in CLBP, including opioid-treated CLBP, and they have shown positive effects on pain intensity, function, and pain coping,[33,36-42] as well as on objectively assessed pain sensitivity to noxious stimuli.[37]

Meditation helps with opioid use and mood effects in people with CLBP as well. Preliminary evidence also suggests that mindfulness-based treatments can assist patients with chronic pain with reducing aberrant opioid-use-related behaviors, desire for opioids, and daily opioid use.[33,43] A systematic review and meta-analysis of mindfulness meditation as a therapy for mental health conditions and well-being showed meditation's efficacy for reducing pain severity in chronic pain conditions as well as decreasing anxiety and depression symptoms.[37] Furthermore, recent systematic reviews found that Mindfulness-Based Therapy can be effective specifically for CLBP.[20,24,44-46]

Relating to Pain in a New Way. Mindfulness-based interventions, including mindfulness meditation practice, facilitate and improve the mind's ability to pay attention to "the present moment" while maintaining an accepting, nonjudgmental awareness of "the present moment" experiences (e.g., thoughts, emotions, or bodily sensations, such as pain).[15,20] As such, meditation may help change the individual's *relationship* to pain and other experiences, rather than focusing on changing the content of the experience itself (which, of note, may not be possible). It has the potential to uncouple the physical experience of pain from pain-related suffering. In chronic pain conditions, pain severity and pain-related suffering have been conceptualized as overlapping but unique entities, further supporting the understanding of chronic pain as a multidimensional construct.[47] Pain coping skills play a major role in how patients cope with and experience pain. Acceptance-based pain coping has been linked to improved outcomes in chronic pain. Mindfulness meditation training may improve adaptive, acceptance-based coping, while reducing unhealthy "fear-avoidance" (e.g., pain catastrophizing) coping.[48,49]

## ELEMENTS OF MINDFUL AWARENESS TRAINING FOR CHRONIC LOW BACK PAIN

Many of the existing mindfulness meditation-based programs have been patterned after Mindfulness-Based Stress Reduction (MBSR[34]), and consist of eight weekly group sessions, guided by experienced, trained instructors, and daily at-home practice. During the program, participants are taught a variety of mindfulness meditation techniques, including longer, more formal techniques (e.g., body scan, breath, or sitting meditation), as well as shorter techniques (e.g., mindful walking, mindfulness of daily activities, other minimeditations) that may be particularly useful for immediate coping with "acute" challenges brought by daily life. A common mini-meditation practice that may be helpful to implement when faced with a challenge such as a pain flare involves the following steps:

- 1. Taking a moment to pause (instead of automatically reacting to the challenge/pain)
- 2. Noticing what is happening "right now" in the body, mind, and heart (the thoughts, images, emotions, and sensations that are occurring)

- 3. Focusing on noticing the sensations of breathing, of the air moving or the belly rising and falling with each breath
- 4. Expanding one's awareness to the body and mind as a whole
- 5. Making a mindful decision about, as opposed to reacting automatically about, what to do next (if anything).[32,33]

#### RESOURCES

For an extensive list of resource materials, refer to "Mindful Awareness," Chapter 4 of the *Passport to Whole Health*.

#### CHRONIC PAIN/CHRONIC LOW BACK PAIN RESOURCES

- 1. Healing and the Mind with Bill Moyers: "Healing from Within" website and video, Bill Moyers (1993)
- 2. Living Well with Pain and Illness: The Mindful Way to Free Yourself from Suffering, Vidyamala Burch (2010)
- 3. Mindfulness Meditation for Pain Relief: Guided Practices for Reclaiming Your Body and Your Life (book, 2-disc CD set), Jon Kabat-Zinn (2009
- 4. Natural Pain Relief: How to Sooth and Dissolve Physical Pain with Mindfulness, Shinzen Young (2011)
- 5. The Mindfulness Solution to Pain: Step-by-Step Techniques for Chronic Pain Management. Jackie Gardner-Nix. (2009)

#### **RESOURCE LINKS**

- Mindful Awareness: https://wholehealth.wisc.edu/overviews/mindful-awareness/
- Power of the Mind: https://wholehealth.wisc.edu/overviews/power-of-the-mind/
- Healing from Within: https://billmoyers.com/series/healing-and-the-mind/

#### **AUTHORS**

"Mindfulness Meditation for Chronic Low Back Pain" was written by <u>Aleksandra E. Zgierska</u>, MD, PhD, and <u>Cindy A. Burzinski</u>, MS, CSAC, LPCT. (2014, updated 2019)

This Whole Health tool (or overview) was made possible through a collaborative effort between the University of Wisconsin Integrative Health Program, VA Office of Patient Centered Care and Cultural Transformation, and Pacific Institute for Research and Evaluation

#### REFERENCES

1. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a

- systematic analysis for the Global Burden of Disease Study 2013. *Lancet.* 2015;386(9995):743-800.
- 2. Martin BI, Turner JA, Mirza SK, Lee MJ, Comstock BA, Deyo RA. Trends in health care expenditures, utilization, and health status among US adults with spine problems, 1997-2006. *Spine*. 2009;34(19):2077-2084.
- 3. Gore M, Sadosky A, Stacey BR, Tai KS, Leslie D. The burden of chronic low back pain: clinical comorbidities, treatment patterns, and health care costs in usual care settings. *Spine*. 2012;37(11):E668-677.
- 4. U.S. Department of Health and Human Services. Healthy People 2020: 2020 Topics and Objectives (Arthritis, Osteoporosis, and Chronic Back Conditions). <a href="http://www.healthypeople.gov/2020/topicsobjectives2020/overview.aspx?topicid=3">http://www.healthypeople.gov/2020/topicsobjectives2020/overview.aspx?topicid=3</a>. Accessed March 31, 2014.
- 5. National Center for Complementary and Integrative Health (NCCIH). NCCIH's New Strategic Plan: Exploring the Science of Complementary and Integrative Health. 2016; NIH website. <a href="https://content.govdelivery.com/accounts/USNIHNCCIH/bulletins/14d3cdd">https://content.govdelivery.com/accounts/USNIHNCCIH/bulletins/14d3cdd</a>. Updated June 3, 2016. Accessed May 5, 2020.
- 6. Meucci RD, Fassa AG, Faria NM. Prevalence of chronic low back pain: systematic review. *Rev Saude Publica*. 2015;49:1.
- 7. Dillie KS, Fleming MF, Mundt MP, French MT. Quality of life associated with daily opioid therapy in a primary care chronic pain sample. *J Am Board Fam Med.* 2008;21(2):108-117.
- 8. Dunn KM, Saunders KW, Rutter CM, et al. Opioid prescriptions for chronic pain and overdose: a cohort study. *Ann Intern Med.* 2010;152(2):85-92.
- 9. Deyo RA, Smith DH, Johnson ES, et al. Opioids for back pain patients: primary care prescribing patterns and use of services. *J Am Board Fam Med.* 2011;24(6):717-727.
- 10. Kobus AM, Smith DH, Morasco BJ, et al. Correlates of higher-dose opioid medication use for low back pain in primary care. *J Pain.* 2012;13(11):1131-1138.
- 11. Koes BW, Backes D, Bindels PJE. Pharmacotherapy for chronic non-specific low back pain: current and future options. *Expert Opin Pharmacother*. 2018;19(6):537-545.
- 12. Tucker HR, Scaff K, McCloud T, et al. Harms and benefits of opioids for management of non-surgical acute and chronic low back pain: a systematic review. *Br J Sports Med.* 2020;54(11):664.
- 13. Gatchel RJ. The continuing and growing epidemic of chronic low back pain. *Healthcare (Basel, Switzerland).* 2015;3(3):838-845.
- 14. Institute of Medicine Committee on Advancing Pain Research C, Education. The National Academies Collection: Reports funded by National Institutes of Health. In: Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research. Washington (DC): National Academies Press (US), National Academy of Sciences.; 2011.
- 15. Bishop SR, Lau M, Shapiro S, et al. Mindfulness: a proposed operational definition. *Clin Psychol Sci.* 2004;11(3):230-241.
- 16. Chiesa A, Serretti A. Mindfulness-based interventions for chronic pain: a systematic review of the evidence. *J Altern Complement Med.* 2011;17(1):83-93.
- 17. Cramer H, Haller H, Lauche R, Dobos G. Mindfulness-based stress reduction for low back pain. a systematic review. *BMC Complement Altern Med.* 2012;12:162.

- 18. Goyal M, Singh S, Sibinga EM, et al. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med.* 2014;174(3):357-368.
- 19. Patil SG. Effectiveness of mindfulness meditation (Vipassana) in the management of chronic low back pain. *Indian J Anaesth.* 2009;53(2):158-163.
- 20. Chou R, Deyo R, Friedly J, et al. Nonpharmacologic therapies for low back pain: a systematic review for an American College of Physicians Clinical Practice Guideline. *Ann Intern Med.* 2017;166(7):493-505.
- 21. Sampaio CV, Lima MG, Ladeia AM. Meditation, health and scientific investigations: review of the literature. *J Relig Health*. 2017;56(2):411-427.
- 22. Demarzo MM, Montero-Marin J, Cuijpers P, et al. The efficacy of mindfulness-based interventions in primary care: A meta-analytic review. *Ann Fam Med.* 2015;13(6):573-582.
- 23. Goldberg SB, Tucker RP, Greene PA, et al. Mindfulness-based interventions for psychiatric disorders: A systematic review and meta-analysis. *Clin Psychol Rev.* 2018;59:52-60.
- 24. Hilton L, Hempel S, Ewing BA, et al. Mindfulness Meditation for chronic pain: systematic review and meta-analysis. *Ann Behav Med.* 2017;51(2):199-213.
- 25. Majeed MH, Ali AA, Sudak DM. Mindfulness-based interventions for chronic pain: Evidence and applications. *Asian J Psychiatr.* 2018;32:79-83.
- 26. Grossman P, Tiefenthaler-Gilmer U, Raysz A, Kesper U. Mindfulness training as an intervention for fibromyalgia: evidence of postintervention and 3-year follow-up benefits in well-being. *Psychother Psychosom.* 2007;76(4):226-233.
- 27. Ball EF, Nur Shafina Muhammad Sharizan E, Franklin G, Rogozinska E. Does mindfulness meditation improve chronic pain? A systematic review. *Curr Opin Obstet Gynecol.* 2017;29(6):359-366.
- 28. Winbush NY, Gross CR, Kreitzer MJ. The effects of mindfulness-based stress reduction on sleep disturbance: a systematic review. *Explore (NY)*. 2007;3(6):585-591.
- 29. Davidson RJ, Kabat-Zinn J, Schumacher J, et al. Alterations in brain and immune function produced by mindfulness meditation. *Psychosom Med.* 2003;65(4):564-570.
- 30. Zeidan F, Vago DR. Mindfulness meditation-based pain relief: a mechanistic account. *Ann N Y Acad Sci.* 2016;1373(1):114-127.
- 31. Segal ZV, Williams JM, Teasdale JD. *Mindfulness-Based Cognitive Therapy for Depression: A New Approach to Preventing Relapse.* New York: Guilford Publications; 2002.
- 32. Bowen S, Chawla N, Marlatt GA. *Mindfulness-based Relapse Prevention for Addictive Behaviors: A Clinician's Guide.* New York, NY: The Guilford Press; 2011.
- 33. Zgierska AE, Burzinski CA, Cox J, et al. Mindfulness Meditation-Based Intervention Is Feasible, Acceptable, and Safe for Chronic Low Back Pain Requiring Long-Term Daily Opioid Therapy. *J Altern Complement Med.* 2016;22(8):610-620.
- 34. Kabat-Zinn J. An outpatient program in behavioral medicine for chronic pain patients based on the practice of mindfulness meditation: theoretical considerations and preliminary results. *Gen Hosp Psychiatry.* 1982;4(1):33-47.
- 35. Kabat-Zinn J, Lipworth L, Burney R. The clinical use of mindfulness meditation for the self-regulation of chronic pain. *J Behav Med.* 1985;8(2):163-190.

- 36. Morone NE, Greco CM, Weiner DK. Mindfulness meditation for the treatment of chronic low back pain in older adults: a randomized controlled pilot study. *Pain.* 2008;134(3):310-319.
- 37. Zgierska AE, Burzinski CA, Cox J, et al. Mindfulness meditation and cognitive behavioral therapy intervention reduces pain severity and sensitivity in opioid-treated chronic low back pain: pilot findings from a randomized controlled trial. *Pain Med.* 2016;17(10):1865-1881.
- 38. Day MA, Ward LC, Ehde DM, et al. A pilot randomized controlled trial comparing Mindfulness Meditation, Cognitive Therapy, and Mindfulness-Based Cognitive Therapy for chronic low back pain. *Pain Med.* 2019;20(11):2134-2148.
- 39. Ardito RB, Pirro PS, Re TS, et al. Mindfulness-Based Stress Reduction program on chronic low-back pain: a study investigating the impact on endocrine, physical, and psychologic functioning. *J Altern Complement Med.* 2017;23(8):615-623.
- 40. Banth S, Ardebil MD. Effectiveness of mindfulness meditation on pain and quality of life of patients with chronic low back pain. *Int J Yoga.* 2015;8(2):128-133.
- 41. Esmer G, Blum J, Rulf J, Pier J. Mindfulness-based stress reduction for failed back surgery syndrome: a randomized controlled trial. *J Am Osteopath Assoc.* 2010;110(11):646-652.
- 42. Garland EL, Thomas E, Howard MO. Mindfulness-oriented recovery enhancement ameliorates the impact of pain on self-reported psychological and physical function among opioid-using chronic pain patients. *J Pain Symptom Manage*. 2014;48(6):1091-1099.
- 43. Garland EL, Manusov EG, Froeliger B, Kelly A, Williams JM, Howard MO. Mindfulness-Oriented Recovery Enhancement for Chronic Pain and Prescription Opioid Misuse: Results From an Early-Stage Randomized Controlled Trial. *J Consult Clin Psychol.* 2014;82(3):448-459.
- 44. Skelly AC, Chou R, Dettori JR, et al. AHRQ Comparative Effectiveness Reviews. In: *Noninvasive Nonpharmacological Treatment for Chronic Pain: A Systematic Review.* Rockville (MD): Agency for Healthcare Research and Quality (US); 2018.
- 45. Jacob JA. As Opioid Prescribing Guidelines Tighten, Mindfulness Meditation Holds Promise for Pain Relief. *JAMA*. 2016;315(22):2385-2387.
- 46. Cherkin DC, Sherman KJ, Balderson BH, et al. Effect of mindfulness-based stress reduction vs cognitive behavioral therapy or usual care on back pain and functional limitations in adults with chronic low back pain: a randomized clinical trial. *JAMA*. 2016;315(12):1240-1249.
- 47. Turk DC, Wilson HD. Pain, suffering, pain-related suffering--are these constructs inextricably linked? *Clin J Pain.* 2009;25(5):353-355.
- 48. de Boer MJ, Steinhagen HE, Versteegen GJ, Struys MM, Sanderman R. Mindfulness, acceptance and catastrophizing in chronic pain. *PLoS One.* 2014;9(1):e87445.
- 49. Schutze R, Rees C, Preece M, Schutze M. Low mindfulness predicts pain catastrophizing in a fear-avoidance model of chronic pain. *Pain.* 2010;148(1):120-127.